



002250-195

Marked-Up Copy of Substitute Specification
Filed January 31, 2005

BOTANICAL COMMERCIAL CLASSIFICATION

Syringa ~~hybrid~~ *meyeri* x *Spyringa microphylla*/Ornamental Lilac Shrub

VARIETAL DENOMINATION

cv. 'Bailming'

SUMMARY OF THE INVENTION

The new Lilac plant of the present invention was created at Harwood, North Dakota, U.S.A., by the cross of *Syringa meyeri* 'Palibin' (non-patented in the United States) and *Syringa microphylla* 'Superba' (non-patented in the United States). The parentage can be summarized as follows:

'Palibin' x 'Superba'.

Seeds from the cross were collected during 1976, were planted, and the resulting plants were observed. A single plant of new cultivar was selected during 1985 in view of its distinctive combination of ornamental characteristics. This plant initially was designated No. 92-1.

It was found that the cultivar of *Syringa meyeri* x *Syringa microphylla* exhibits the following combination of characteristics:

- (a) exhibits a rounded uniform compact growth habit,

Marked-Up Copy of Substitute Specification
Filed January 31, 2005

- (b) forms attractive trusses of fragrant red purple blossoms in late spring that fade to lavender pink,
- (c) is relatively free of insect and disease problems, and
- (d) is well-suited for growing as a distinctive ornamental shrub in the landscape.

The new cultivar of the present invention provides a distinctive red purple blossom coloration to the landscape that is displayed in late spring after many other shrubs have finished flowering. It can be grown to advantage as a specimen shrub or in a mass planting. The plant is versatile for many uses in the landscape including foundation plantings and informal hedges. The superior hardiness of the *Syringa meyeri* 'Palibin' parent which generally is lacking in the *Syringa microphylla* 'Superba' parent is imparted to the new cultivar of the present invention.

The new cultivar of the present invention can be readily distinguished from its parental cultivars. More specifically, the 'Palibin' cultivar exhibits a low spreading growth habit and the blossoms open to a pale lilac coloration, and the 'Superba' cultivar exhibits a taller and considerably more spreading growth habit and forms deep pink blossoms.

Marked-Up Copy of Substitute Specification
Filed January 31, 2005

The new cultivar of the present invention also can be readily distinguished from other Lilac cultivars, such as 'Bailbelle' (United States Plant Patent No. 12,294), 'Baildust' (United States Plant Patent ~~Application No. 10/158,047,~~ ~~filed May 31, 2002~~) (No. 15,152), and 'Bailsugar' (United States Plant Patent Application No. 10/365,435, filed February 13, 2003), resulting from the same breeding program in view of the distinctive blossom coloration. More specifically, the 'Bailbelle' variety forms pink blossoms, the 'Baildust' variety forms pale antique pink blossoms, and the 'Bailsugar' variety forms purple lilac blossoms. The red purple blossoms of the new variety fade to an attractive lavender pink.

The new cultivar of the present invention was asexually propagated by the use of rooted cuttings and grafting at St. Paul, Minnesota, U.S.A., during 1992, and the progeny were field planted during 1993. The distinctive characteristics of the new cultivar have been found to be stable and to be capable of transmission from one generation to another following such asexual propagation at St. Paul, Minnesota and elsewhere. The new cultivar reproduces true to type in successive generations of asexual reproduction.

Marked-Up Copy of Substitute Specification
Filed January 31, 2005

The new cultivar of the present invention also has been grafted on *Syringa reticulata* to form distinctive small grafted trees.

The new cultivar of the present invention has been named 'Bailming'. It is a member of the FAIRYTALE™ Series of Lilac plants that includes the 'Bailbelle' 'Baildust' and 'Bailsugar' cultivars, and is being marketed under the PRINCE CHARMING trademark.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as true as reasonably possible to make the same in color illustrations of this character typical plants of the new cultivar. The plants were propagated by use of rooted cuttings and were photographed when approximately three years of age while growing outdoors during late spring at St. Paul, Minnesota, U.S.A.

FIG. 1 - shows a blossoming plant wherein the uniformly rounded and compact growth habit of the new cultivar is exhibited.

FIG. 2 - shows a closer view of the red purple blossoms that fade to lavender pink as well as the foliage.

FIG. 3 - shows the distinctive wine red buds with attractive foliage.

Marked-Up Copy of Substitute Specification
Filed January 31, 2005

FIG. 4 - shows a close-up view of the open blossoms with some foliage.

DETAILED DESCRIPTION

The chart used in the identification of colors is the R.H.S. Colour Chart of the Royal Horticultural Society, London, England. Common terms are to be accorded their ordinary dictionary significance. The description is based upon the observation during July 2002 and 2003 of five year-old plants propagated by the use of rooted cuttings and growing in containers and in the field at St. Paul, Minnesota, U.S.A.

BOTANICAL CLASSIFICATION: *Syringa meyeri* x *Syringa microphylla*,
cv. 'Bailming'.

PLANT:

Growth habit. -- Neat compact rounded shrub, and a
uniform grower.

Marked-Up Copy of Substitute Specification
Filed January 31, 2005

Size. -- Forms a shrub of approximately 1.5 to
1.8 m in height and approximately 1.25 to
1.8 m width.

BARK:

Texture. -- Glabrous.
Color. -- Grey-Brown Group 199B.

FOLIAGE:

Leaf shape. -- Elliptic/ovate.
Arrangement. -- Opposite.
Leaf apex. -- Acute to obtuse.
Leaf base. -- Rounded.
Leaf surface. -- Matte.
Leaf margins. -- Entire.
Venation. -- Palmately-veined.
Color. -- Yellow-Green Group 146A blending with
Red-Purple Group 59B at the margin
(upper surface) and Yellow-Green Group

Marked-Up Copy of Substitute Specification
Filed January 31, 2005

146B suffused with Red-Purple Group 59A (under surface) for young foliage, and Yellow-Green Group 147A (upper surface) and Yellow-Green Group 147B and 147C with veins of Yellow-Green Group 147A (under surface) for adult foliage. This can be compared to Green Group 139A (upper surface) for the adult foliage of the *Syringa meyeri* 'Palibin' parent, and Green Group 137A (upper surface) and Green Group 138C (under surface) for the adult foliage of the 'Bailbelle' cultivar.

Petioles. -- Yellow-Green Group 147C on the upper surface and Yellow-Green Group 147C blending with Red-Purple Group 59A at the axis on the under surface.

Stems. -- Yellow-Green Group 146B on young stems and Greyed-Green Group 197A on adult wood.

Marked-Up Copy of Substitute Specification
Filed January 31, 2005

Lenticels. -- Commonly are present on stems and
Grey-Brown Group 199D in coloration.

INFLORESCENCE

~~*Arrangement.*~~ -- ~~Opposite.~~

Age to Flower. == Typically the first flowers appear the second year from a rooted cutting in the field and in container production. A few flowers have been observed during the first year sometimes by late summer depending upon the time of planting.

Time of Flowering. == Typically mid- to late-May at the indicated location, and commonly after the 'Bailbelle' cultivar. Such flowering commonly is after many other flowering shrubs have finished blooming.

Marked-Up Copy of Substitute Specification
Filed January 31, 2005

- Buds.* -- In trusses, closed funnel-form, red purple, Red-Purple Group 70A, (upper surface) and Red-Purple Group 71B (under surface). The length commonly is 1.5 cm and width is approximately 0.4 cm on average.
- Flower Arrangement.* -- Densely packed in panicles approximately 8 to 9 cm in length and approximately 4 to 5 cm in width.
- Flower Configuration.* -- Tubular with four petal lobes at the end (as illustrated).
- Flower Size.* -- Approximately 1.0 cm in length and approximately 0.65 cm in diameter at the widest point.
- Flower Color.* -- When first opening the petals are red purple, Red-Purple Group 70A, with Red-Purple Group 69A at the margin (upper surface) and Red-Purple Group 70B with Red-Purple Group 69B and 69C at the

Marked-Up Copy of Substitute Specification
Filed January 31, 2005

margin (under surface). When blooming the petals are Red-Purple Group 70C with Red-Purple Group 69A at the margin (upper surface) and red purple, Red-Purple Group 70B with Red-Purple Group 69B and 69C at the margin (under surface). At the end of blooming the petals are Red-Purple Group 75A (upper surface) and Red-Purple Group 74D (under surface).

Stamens. -- Two per floret, and near Red-Purple Group 68A with some near White Group 155D in coloration.

Filaments. -- Red-Purple Group 62D in coloration.

Anthers. -- Red-Purple Group 64D in coloration.

Pollen. -- Present.

Pistil. -- Approximately 0.4 cm in length.

Style. -- Red-Purple Group 62C in coloration.

HARDINESS

-11-

Marked-Up Copy of Substitute Specification
Filed January 31, 2005

CULTURE

Similar to the Dwarf Korean Lilac. Prefers well-drained soil and a sunny growing location. Generally does not well tolerate poorly drained growing conditions.

DISEASE/PEST RESISTANCE

Has proven to be relatively free of disease and insect problems during observations to date.

LANDSCAPE USAGE

Provides a hardy highly ornamental fragrant shrub having a distinctive blossom coloration that can be grown as a specimen plant or in a mass planting. Can be grown as a foundation planting or as an informal hedge.